

Convex relaxations of mixed derivatives of multivariate entire functions/polynomials

Leonid Gurvits *

Abstract

Many "hard" combinatorial and geometric quantities (such as the number of perfect matchings in bipartite graphs (the **permanent**), number of perfect matchings in general graphs (the **hafnian**), the number of matching of the fixed size, the number of Hamiltonian cycles, the number of exact 3-coverings, the number of common bases in the intersection of unimodular geometric matroid and the matroid of transversals (the **Mixed Discriminant**), the **Mixed Volume** (responsible for the number of isolated solutions of systems of polynomial equations) etc.) can be expressed as the coefficient $a_{1,\dots,1} = \frac{\partial^n}{\partial x_1 \dots \partial x_n} p(0, 0, \dots, 0)$ of the monomial $x_1 \dots x_n$ in some effectively computable "generating" homogeneous polynomial $p \in Hom_+(n, n)$ of degree n with nonnegative rational coefficients. In other words the counting problems are particular cases of partial (for we are after only one coefficient) multivariate interpolations; and the corresponding decision problems (existence of perfect matchings, existence of Hamiltonian cycles etc.) are equivalent to checking if " $\frac{\partial^n}{\partial x_1 \dots \partial x_n} p(0, 0, \dots, 0) = 0$?". Define the next quantity, associated with a homogeneous polynomial $p \in Hom_+(n, n)$:

$$Cap(p) =: \inf_{x_i > 0} \frac{p(x_1, \dots, x_n)}{\prod_{1 \leq i \leq n} x_i}.$$

Note that $\log(Cap(p)) = \inf_{y_1 + \dots + y_n = 0} \log(p(e^{y_1}, \dots, e^{y_n}))$ and the functional $\log(p(e^{y_1}, \dots, e^{y_n}))$ is convex. (*I will explain that the monotonicity of the celebrated Baum-Welsh algorithm for HMM is due to the above convexity.*)

Therefore $\log(Cap(p))$ can be, with some extra care and luck, effectively additively approximated using convex programming tools and an oracle, deterministic or random, evaluating the polynomial p . It directly follows from the nonnegativity of the coefficients of p that

$$\log(Cap(p)) \geq \log\left(\frac{\partial^n}{\partial x_1 \dots \partial x_n} p(0, 0, \dots, 0)\right).$$

In order to show that $\log(Cap(p))$ is a reasonable relaxation for $\log\left(\frac{\partial^n}{\partial x_1 \dots \partial x_n} p(0, 0, \dots, 0)\right)$ it is desirable to prove the (optimal) reverse bound:

$$\frac{\partial^n}{\partial x_1 \dots \partial x_n} p(0, 0, \dots, 0) \geq Const(n, p) Cap(p), Const(n, p) > 0.$$

Such lower bound is impossible in general, but I will describe a rather broad class of polynomials (and even of entire functions) for which it is possible.

The combinatorial ([1],[2]), [4], [3], algorithmic ([5]) and "structural" [6] applications will be described. Most importantly, a fairly complete proof will be presented.

*gurvits@lanl.gov. Los Alamos National Laboratory, Los Alamos, NM.

References

- [1] A. Schrijver, Counting 1-factors in regular bipartite graphs, *Journal of Combinatorial Theory, Series B* 72 (1998) 122–135.
- [2] M. Laurent and A. Schrijver, On Leonid Gurvits' proof for permanents, 2009, <http://homepages.cwi.nl/~lex/files/perma5.pdf>, to appear in *American Mathematical Monthly*.
- [3] S. Friedland and L. Gurvits, Lower Bounds for Partial Matchings in Regular Bipartite Graphs and Applications to the Monomer-Dimer Entropy, *Combinatorics, Probability and Computing*, 2008.
- [4] L. Gurvits, Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all, *Electronic Journal of Combinatorics* 15 (2008).
- [5] L. Gurvits, A polynomial-time algorithm to approximate the mixed volume within a simply exponential factor. *Discrete Comput. Geom.* 41 (2009), no. 4, 533–555. On multivariate Newton-like inequalities
- [6] L. Gurvits, On multivariate Newton-like inequalities, <http://arxiv.org/abs/0812.3687>.